



UNIVERSITI PUTRA MALAYSIA

MICROPROPAGATION OF *MICHELIA CHAMPACA* L.

**ARMİYANTI
FP 2009 25**

MICROPROPAGATION OF *MICHELIA CHAMPACA* L.

ARMIYANTI

**MASTER OF SCIENCE
UNIVERSITI PUTRA MALAYSIA
2009**



MICROPROPAGATION OF *MICHELIA CHAMPACA* L.

By
ARMİYANTI

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

October 2009



Specially Dedicated

To

**My Late Father and Late Mother:
Ansari Itam and Arifah**

*..... who inspired, supported and gave me tremendous
courage to be a good person*

All my relatives and my beloved husband

*..... who inspired, supported, patience and understanding
during the period of my study*

Abstract of thesis presented to the Senate of Universiti Putra Malaysia
in fulfilment of the requirement for the degree of Master of Science

MICROPROPAGATION OF *MICHELIA CHAMPACA* L.

By

ARMİYANTI

October 2009

Chairman : Mihdzar Abdul Kadir, PhD

Faculty : Agriculture

The expansion in Champaca industry has led to an increasing demand for planting materials. A study was conducted with the objective of developing plant regeneration system for *Michelia champaca* L. through organogenesis and somatic embryogenesis. The study was conducted in consideration of the potential economic importance of the species. It consisted of development of sterilization procedure for field grown *M. champaca* shoot tip and nodal segment explants and plant regeneration through organogenesis and somatic embryogenesis of *M. champaca* either through solid culture medium and cell suspension culture technique. This was the first report on the establishment of *M. champaca* cell suspension culture.

All experiments in this study were conducted in a Completely Randomized Design (CRD). In the development of sterilization procedure for field grown shoot tips and nodal segments of *M. champaca* sterilization with 20% of clorox (20 minutes) + 70% alcohol (2 minutes) + 10% clorox (5 minutes) and 0.1% HgCl₂ (5 minutes)

successfully produced non-contaminated explants up to 80% for field-grown shoot tip and 50% for nodal explants.

In the experiment of determination of *M. champaca* seed viability, different tetrazolium concentrations (0.1, 0.5, and 1 % (w/v)) in combination with different immersion times (1, 2, 4, 6 and 8 hours) were used as treatments. The highest percentage of seed viability (100%) was obtained from all tetrazolium concentrations tested in combination with eight hours of immersion time.

Results on plant regeneration via organogenesis showed that MS (Murashige and Skoog, 1962) medium supplemented with 0.2 gL⁻¹ of charcoal, 30 gL⁻¹ (w/v) of sucrose and solidified with 3.9 gL⁻¹ gelrite agar containing BAP at 6 mgL⁻¹ in combination with 0.5 mgL⁻¹NAA was the most suitable for shoot induction from shoot tip derived from seedling explants with the highest mean number of shoots produced per explant (1.8) and mean shoot height of 4.53 cm. Meanwhile treatment containing 0.5 mgL⁻¹ NAA was the most suitable for rooting with a percentage of shoot producing root at 70%, with mean number of roots formed per shoot at 2.5 and mean root length was 1.07 cm.

For plant regeneration through somatic embryogenesis using immature seed, the highest percentage of explants that responded to form somatic embryos (30%) and mean number of somatic embryos produced per explant (87.3) were obtained on treatment containing 2 mgL⁻¹ NAA. However, higher frequency of somatic embryo

formation was achieved through cell suspension culture system. Liquid full strength MS medium supplemented with 2 mgL⁻¹ (w/v) NAA produced a mean number 47.67±4.53 per ml cotyledonary somatic embryos.

For germination of somatic embryos produced from solid medium and through cell suspension culture, hormone free MS medium gave the highest percentage of somatic embryo germination at 56% and 34%, and normal plantlet production at 45% and 29% respectively.

Plantlet regeneration protocols through organogenesis and somatic embryogenesis of *M. champaca* which were successfully established in this study could be used for mass production of planting materials of this crop.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

PEMBIAKAN MIKRO *MICHELIA CHAMPACA* L.

Oleh

ARMİYANTI

Oktober 2009

Pengerusi : Mihdzar Abdul Kadir, PhD

Fakulti : Pertanian

Perluasan industri Champaca menyebabkan peningkatan permintaan kepada bahan tanaman. Suatu kajian telah dijalankan dengan matlamat untuk mengembangkan kaedah regenerasi tumbuhan keatas *Michelia champaca* L. melalui organogenesis dan embriogenesis soma. Kajian ini dijalankan berdasarkan potensi kepentingan ekonomi spesies ini. Kajian ini terdiri daripada pembinaan kaedah pensterilan keatas eksplan mercu pucuk dan keratan ruas *M. champaca* yang ditanam di lapangan dan regenerasi tumbuhan melalui organogenesis dan embriogenesis soma samada melalui media kultur pepejal dan teknik kultur ampai sel. Kajian ini adalah laporan pertama seumpama terhadap penentuan kultur ampai sel daripada *M. champaca*.

Kesemua kajian ini disusun dalam reka bentuk berawak penuh (CRD). Dalam pembinaan kaedah pensterilan keatas eksplan mercu pucuk dan keratan ruas *M. champaca* yang ditanam di lapangan, kaedah pensterilan dengan clorox 20% (20 minit) + alkohol 70% (2 minit) + clorox 10% (5 minit) dan HgCl₂ 0.1% (5 minit)

telah berjaya menghasilkan eksplan mercu pucuk yang ditanam di lapangan terbebas daripada kontaminasi sebanyak 80% dan 50% keatas eksplan buku ruas.

Pada kajian penentuan viabiliti biji *M. champaca*, kepekatan tetrazolium yang berbeza (0.1, 0.5, dan 1 % (b/v)) yang dikombinasikan dengan masa perendaman yang berbeza (1, 2, 4, 6 dan 8 jam) telah digunakan sebagai rawatan. Peratusan viabiliti biji tertinggi (100%) diperolehi dari keseluruhan kepekatan tetrazolium yang dicuba dengan kombinasi lapan jam perendaman.

Hasil keatas regenerasi tumbuhan melalui organogenesis telah menunjukkan bahawa media MS (Murashige and Skoog, 1962) yang ditambahkan 0.2 gL^{-1} arang, 30 gL^{-1} (b/v) sukrosa dan dipejalkan dengan 3.9 gL^{-1} agar gelrite dan mengandungi BAP 6 mgL^{-1} dikombinasikan dengan 0.5 mgL^{-1} NAA adalah yang paling sesuai untuk induksi tunas daripada eksplan mercu pucuk yang diperolehi daripada anak benih dengan purata bilangan tunas terhasil yang paling tinggi per eksplan (1.8) dan purata tinggi tunas sebesar 4.53 cm. Manakala, untuk pengakaran, rawatan yang mengandungi 0.5 mgL^{-1} NAA adalah paling sesuai untuk pengakaran dengan peratusan tunas yang menghasilkan akar sebanyak 70% dengan purata bilangan akar terbentuk setiap tunas adalah 2.5 dan purata panjang akar adalah 1.07 cm.

Untuk regenerasi tumbuhan melalui embriogenesis soma dengan menggunakan biji muda, peratusan paling tinggi daripada eksplan membentuk embrio soma (30%) dan purata bilangan embrio soma yang terhasil per eksplan (87.3) telah diperolehi

daripada rawatan yang mengandungi 2 mgL^{-1} NAA. Walau bagaimanapun, frekuensi pembentukan embrio soma yang lebih tinggi telah dicapai melalui kaedah kultur ampai sel. Medium penuh cecair MS yang mengandungi 2 mgL^{-1} (b/v) NAA telah menghasilkan kotiledon embrio soma sebanyak 47.67 ± 4.53 per ml.

Untuk percambahan embrio soma yang dihasilkan daripada kaedah media pejal dan kaedah kultur ampai sel, medium MS yang tidak mengandungi hormon telah memberikan peratusan embrio soma bercambah paling tinggi masing-masing pada 56% dan 34% dan penghasilan anak pokok normal masing-masing pada 45% dan 29%.

Kaedah regenerasi anak pokok melalui organogenesis dan embriogenesis soma daripada *M. champaca* yang telah berjaya dilaksanakan dalam kajian ini dapat digunakan untuk pengeluaran besar-besaran bahan tanaman daripada tumbuhan ini.

ACKNOWLEDGEMENTS

In the name of Allah The Beneficial and The Compassionate

Praise be to Allah the Most Gracious and Merciful, upon His permission I could complete this thesis well.

I would like to express my sincere thanks to my supervisor, Associate Professor Dr. Mihdzar Abdul Kadir of the Department of Agriculture Technology, Faculty of Agriculture for his supervision, kind support beyond duties and at heart for being so patient and understanding.

I am also indebted to my co-supervisor, Associate Professor Dr Saleh Kadzimin, Department of Crop Science, Faculty of Agriculture for his guidance and constructive suggestion during the course of this project.

I would also like to express my deepest thanks to my late father Ansari Itam, my late mother Arifah, my beloved husband Syaiful Bahri Panjaitan, my elder sister Arlida, Arlina, Arlaini, Anizar, Ardiana, my brother Afrul Wahyuni and my youngest sister Afriani, all my brother in law (Ulli, Azmi, Hermanto, Muhammad Bahi, and Zainal), my niece and nephew (Aldi, Reza, Intan, Ula, Fiqhan, Amirul, Ilham, Syaukal, Pasya and Aisyah), my best friend Ria, Aswani, Naimah, Susi, Kak Raudah, Melly and Kak Aziah for their encouragement, patience, moral support and inspiration during the period of my study.

I would like to express my sincere gratitude to the State Government of Nanggroe Aceh Darussalam (Indonesia) and the Ministry of Higher Education Malaysia for the scholarships given to me to accomplish this goal.

Finally, special thanks to my lab mates Ibu Innaka, Rozila, Siti Raziah, Shikin, Chia, Fatemeh, Hossein Kamaladini, Azadeh, Motaharreh, Nurul Azwana, Ashida, Maryam, Hasnul, Hanif and Ali Reza Koshravi for their cooperation during the period of my study.

I certify that a Thesis Examination Committee has met on 9th October 2009 to conduct the final examination of Armiyanti on her thesis entitled “Micropropagation of *Michelia champaca* L.” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Pertanian Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Mahmud Tengku Muda Mohammed, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Chairperson)

Maheran Abdul Aziz, PhD

Associate Professor
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Farida Kamaruzzaman, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Ahamad Tarmizi Hashim, PhD

Senior Research Officer
Malaysia Palm Oil Board (MPOB)
43000 Kajang, Selangor Darul Ehsan,
Malaysia
(External Examiner)

BUJANG KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies,
Universiti Putra Malaysia

Date : 24 December 2009

The thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Master of Science. Members of the Supervisory Committee were as follows:

Mihdzar Abdul Kadir, PhD

Associate Professor
Department of Agriculture Technology,
Faculty of Agriculture,
Universiti Putra Malaysia.
(Chairman)

Saleh Kadzimin, Ph.D

Associate Professor
Department of Crop Science,
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date : 14 January 2010

DECLARATION

I hereby declare that this thesis is based on my original work except quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.

ARMİYANTI

Date : 9th October 2009

TABLE OF CONTENTS

DEDICATION	Page ii
ABSTRACT	iii
ACKNOWLEDGEMENT	ix
APPROVAL	xi
DECLARATION	xiii
LIST OF TABLES	xvii
LIST OF FIGURES	xix
LIST OF ABBREVIATIONS	xxv
CHAPTER	
1. INTRODUCTION	
1.1. Background	1.1
1.2. Objectives	1.4
1.3. Significance of Study	1.4
2. LITERATURE REVIEW	
2.1. Botany of Champaca	2.1
2.2. Uses and Commercial Importance of <i>Michelia champaca</i>	2.2
2.3. Champaca Propagation	2.3
2.3.1. Seed propagation	2.3
2.3.2. Vegetative propagation	2.5
2.4. <i>In-Vitro</i> Plant Propagation Systems	2.6
2.4.1. Shoot tip culture from field-grown plants	2.7
2.4.2. Organogenesis	2.8
2.4.3. Somatic embryogenesis	2.9
2.5. Controlling factors in Micropropagation	2.10
2.5.1. Plant growth regulators (PGRs)	2.11
2.5.2. Explant types	2.16
2.5.3. Medium used	2.16
2.6. Sterilisation Techniques in Plant Micropropagation	2.17
2.6.1. Microbial contamination in tissue culture	2.18
2.6.2. Explant surface sterilization	2.19
3. THE DEVELOPMENT OF STERILIZATION PROCEDURES FOR FIELD GROWN SHOOT TIPS AND NODAL EXPLANTS OF <i>MICHELIA CHAMPACA</i>	
3.1. Preface	3.1
3.2. Materials and Methods	3.2
3.3. Results and Discussions	3.5
3.3. Conclusion	3.11



4. IN-VITRO PLANTLET REGENERATION OF <i>MICHELIA CHAMPACA</i> THROUGH ORGANOGENESIS	
4.1. Preface	4.1
4.2. Materials and Methods	4.3
4.3. Results and Discussions	4.15
4.3.1. Determination of <i>Michelia champaca</i> seed viability using tetrazolium stain solution	4.15
4.3.2. Induction of shoots from field-grown shoot tip explant	4.18
4.3.3. Induction of shoot from shoot tip derived from seedlings of <i>Champaca</i>	4.23
4.3.4. Induction of root on shoots regenerated from seedling explants	4.24
4.4. Conclusion	4.31
5. IN-VITRO PLANTLET REGENERATION OF <i>MICHELIA CHAMPACA</i> THROUGH SOMATIC EMBRYOGENESIS	
5.1. Preface	5.1
5.2. Materials and Methods	5.2
5.3. Results and Discussions	5.12
5.3.1. Induction of somatic embryo from immature seeds of <i>M. champaca</i>	5.12
5.3.2. Establishment of cell suspension culture	5.26
5.3.3. Germination of somatic embryos obtained from solid culture medium and cell suspension culture techniques	5.37
5.4. Conclusion	5.47
6. GENERAL DISCUSSION	
6.1. The Development of Sterilization Procedure for Field-Grown <i>Michelia champaca</i> Shoot Tip and Nodal Segment Explants	6.1
6.2. <i>In vitro</i> Plantlet Regeneration of <i>Michelia champaca</i> Through Organogenesis	6.2
6.3. <i>In Vitro</i> Plantlet Regeneration of <i>Michelia champaca</i> Through Somatic Embryogenesis	6.5
7. GENERAL CONCLUSION	
7.1. Conclusions	7.1
7.2. Future Direction	7.2
REFERENCES	
APPENDICES	
BIODATA OF STUDENT	

LIST OF TABLES

Table	Page
3.1. The effect of different surface sterilization treatments on percentage of survival and contamination of field-grown shoot tips of Champaca after one month culture	3.8
3.2. The effect of different surface sterilization treatments on percentage of survival and contamination of nodal segment explants	3.9
4.1. Different Tetrazolium concentrations and time of immersion seed viability (%) of <i>M. champaca</i>	4.6
4.2. The different Cytokinin types (BAP and Kinetin) and concentrations either alone or in combination with 0-0.5 mgL ⁻¹ NAA for shoot induction from field-grown shoot tip and shoot tip derived from seedling of Champaca	4.10
4.3. The effects of cytokinin types (BAP and Kinetin) either alone or in combination with 0.5 mgL ⁻¹ NAA on the percentage of field-grown shoot tip of Champaca producing shoot and the mean number of shoots produced per explant after eight weeks of culture	4.21
4.4. Effects of cytokinin (BAP and Kinetin) either alone or in combination with 0.5 mgL ⁻¹ NAA on the percentage of seedlings derived shoot tip of Champaca producing shoot, the mean number of shoots produced per explant and mean of shoot height after the fourth subculture	4.29
4.5. The effects of auxin types (NAA and IBA) and their concentrations on the percentage of root formation on shoots regenerated from shoot tip explants derived from Champaca seedlings, showing mean number of roots formed per explant and mean root length (cm) after ten weeks of culture	4.32
5.1. Different concentrations of GA3 for germination of somatic embryos from immature seed of <i>M. champaca</i>	5.11
5.2. The estimated yield of somatic embryos in cell suspension cultures established from immature seed of Champaca explants in full MS medium containing 2 mgL ⁻¹ NAA	5.36

- 5.3. Effect of different concentrations of GA3 on the percentage of germination of somatic embryos produced through solid culture medium, the percentage of normal plantlet produced and percentage of abnormal plantlets produced after 12 weeks of culture 5.44
- 5.4. Effect of different concentrations of GA3 on the percentage of germination of somatic embryos produced through cell suspension culture, the percentage of normal plantlet produced and percentage of abnormal plantlets produced after 12 weeks of culture 5.45

LIST OF FIGURES

Figure	Page
1.1. <i>Michelia champaca</i> tree Bar=30cm (A) and flower of <i>M. champaca</i> , Bar=10mm (B)	1.1
3.1. Shoot tip (A) and nodal (B) explants excised to 1.5 cm before culture onto culture medium	3.4
3.2. Contaminated nodal segment explant caused by bacterial (red arrow) (A); Contaminated field-grown shoot tip of champaca explant caused by fungus (blue arrow) (B); Contaminated explants died after one month of culture (C) (Bar=1cm)	3.6
3.3. The field-grown shoot tip of champaca explant (A), and nodal segment explant survived (B) after one month of culture through surface sterilization protocol (T5), (Bar=1cm)	3.7
4.1. Agregate fruit of <i>M. champaca</i> (A); Seeds inside the carpel (B); Seeds of <i>M. champaca</i> after isolate from seed carpel (C) (Bar=1cm)	4.4
4.2. The mature seeds of <i>M. champaca</i> were immersed in distilled water for two days (A); Champaca seeds after removing seed pulp (B); Champaca seeds after removing seed coat (C) (Bar=5mm)	4.5
4.3. Shoot tip explants harvested from field-grown Champaca plant (A) Field-grown shoot tip explants after removal of outer cover of the shoot tip and ready to be cultured (B) (Bar=5mm)	4.9
4.4. Shoot regenerated from shoot tip explant of seedling with old leaves removed and cultured on hormone free MS medium for one week (Bar=4mm)	4.12
4.5. After one week on hormone free MS medium, shoots regenerated from shoot tip explants derived from seedlings were cultured on a rooting medium containing different auxin types (IBA and NAA) and their concentrations (Bar = 4mm)	4.13

4.6.	Viable seeds of Champaca with more than 50% of seed surface stained red (Bar=1mm)	4.15
4.7.	The non viable seeds of Champaca with less than 50% of seed surface stained red (Bar=1mm)	4.16
4.8.	Effect of different Tetrazolium concentrations and times of immersion on <i>M. champaca</i> seed viability (%). Vertical bars represent \pm SD. Bars followed by the same letter(s) are not significantly different using DNMR at $p=0.05$	4.16
4.9.	Field-grown shoot tip of Champaca explant showing senescence at leaf tips (blue arrow) after one week of culture (A); The young leaves opened up and grew (white arrow) after two weeks of culture (B) (Bar= 5mm)	4.18
4.10.	Field-grown shoot tip explant enlarged with axillary shoot primordia formed at the base of explant (white arrow) after four weeks of culture (A); Axillary shoot (white arrow) and shoot tip (blue arrow) elongated at the fifth week of culture (B); The axillary shoot produced turned brown (white arrow) after seven weeks of culture (C); Regenerated finally died (blue arrow) at the eighth weeks of culture (D) (Bar= 5mm)	4.20
4.11.	Shoot tip explant isolated from Champaca seedlings cultured on MS medium containing different concentrations and cytokinin types (BAP and Kinetin) either alone or combined with 0.5 mgL^{-1} of NAA for shoot induction (A); The shoot tip explant showed their response by enlarging and epicotyl started to elongate (white arrow) within two weeks of culture (B); The shoot tip (white arrow) grew and the first and the second young leaves (blue arrow) opened up after four weeks of culture (C); The axillary shoots primordia (blue arrow) grew from between the leaf and stem and shoot tip (white arrow) were then started to grow up further at the sixth week of culture (D). (Bar= 5mm)	4.25
4.12.	The developmental stages of shoot formation from shoot tip derived from seedlings of champaca explants. The shoot tip explant with small axillary shoot (white arrow) grew from between stem ten weeks of culture (A); At the thirteenth week of culture, axillary shoot elongates (white arrow) (B); Axillary shoot proliferation (white arrow) obtained after 18 th week of culture (two weeks after the fourth subculture) on MS medium contained 6 mgL^{-1} BAP in combination with 0.5 mgL^{-1} NAA (C). (Bar= 5mm)	4.26

4.13.	Root primordia (white arrow) formed after three weeks of culture (Bar= 1cm)	4.34
4.14.	Root formation from micro shoots regenerated from seedlings derived shoot tip of champaca on medium containing 0.5 mgL ⁻¹ NAA after ten weeks of culture (Bar= 1cm)	4.35
4.15.	Root formation from micro shoots regenerated from seedlings derived shoot tip of Champaca on rooting medium containing 1 mgL ⁻¹ IBA after ten weeks of culture (Bar= 1cm)	4.35
5.1.	Immature seeds of <i>M. champaca</i> before removing from fruit pericarp (red arrow) (Bar=1cm)	5.4
5.2.	Immature seed of <i>M. champaca</i> with cut at the corner side of seed (red arrow) cultured on somatic embryos induction medium (Bar=2mm)	5.5
5.3.	The explants started to form callus at the 4 th week of culture on all 2,4-D treatments (A). Meanwhile, the explants started to form callus at the 6 th week of culture on all NAA treatments (B). (Bar=5mm)	5.13
5.4.	Callus formation from immature seed of <i>M. champaca</i> on MS medium containing 10 mgL ⁻¹ 2,4-D on the 3 rd month of cultured (Bar=5mm)	5.14
5.5.	Callus formation from immature seed of <i>M. champaca</i> on MS medium containing 8 mgL ⁻¹ NAA at the 3 rd month of culture (Bar=5mm)	5.14
5.6.	Effect of different auxin types (2,4-D and NAA) and their concentrations on the percentage of callus formation from immature seed of <i>M. champaca</i> after six months of culture. Vertical bars represent \pm SD. Bars followed by the same letter(s) are not significantly different using DNMRT at p=0.05	5.15
5.7.	Effect of different auxin types (2,4-D and NAA) and their concentrations on the percentage of embryogenic callus formation from immature seed of <i>M. champaca</i> after six months of culture. Vertical bars represent \pm SD. Bars followed by the same letter(s) are not significantly different using DNMRT at p=0.05	5.18
5.8.	Embryogenic callus from immature seed of Champaca (2 mg/l NAA 4 th month after cultured) (Bar=2mm)	5.20

5.9.	Somatic embryos at heart stage (red arrow) produced from immature seed of <i>M. champaca</i> on MS medium containing 2mgL^{-1} after the 5 th month of culture (Bar=1mm)	5.22
5.10.	Somatic embryos at torpedo stage (red arrow) produced from immature seed of <i>M. champaca</i> on MS medium containing 2mgL^{-1} NAA after the 6 th month of culture (Bar=2mm)	5.22
5.11.	Effect of different auxin types (2,4-D and NAA) and their concentrations on the percentage of explants formed somatic embryos after six months of culture. Vertical bars represent \pm SD. Bars followed by the same letter(s) are not significantly different using DNMR at $p=0.05$	5.23
5.12.	Effect of different auxin types (2,4-D and NAA) and their concentrations on mean number of somatic embryos produced per immature seeds of <i>M. champaca</i> explants after six months of culture. Vertical bars represent \pm SD. Bars followed by the same letter(s) are not significantly different using DNMR at $p=0.05$	5.25
5.13.	Cell growths in liquid MS medium containing 2mgL^{-1} NAA. Vertical bars represent \pm SD (n=10)	5.27
5.14.	Diagram of a standard haemocytometer chamber	5.29
5.15.	The number of cells were counted on haemocytometer	5.29
5.16.	Rounded to elongate and oval shaped cells with varied cell size and cell content (cytoplasm and vacuola) during cell suspension establishment (Bar= $10\mu\text{m}$)	5.30
5.17.	Viable cells with dense cytoplasm fluoresced apple green when assessed using FDA solution (Bar = $20\mu\text{m}$)	5.30
5.18.	Turbidity of cell suspension indicating more cell becoming embryogenic after three to four subcultures (Bar=1cm)	5.32
5.19.	Single cell undergoing to divide into two (A) and (B), the cell divided into four (C), and the cell divided into eight (D), (Bar= $10\mu\text{m}$)	5.32
5.20.	Pro-embryogenic masses in liquid cell suspension culture at the second subculture (A) and after the third subculture (B), (Bar= $10\mu\text{m}$)	5.33

5.21.	Globular somatic embryos in liquid cell suspension culture (A) and single globular somatic embryo formation in liquid cell suspension culture (B) observed under an inverted microscope (Bar=50µm)	5.33
5.22.	Heart stages of somatic embryos in liquid cell suspension culture (A); Early stages of heart shape somatic embryo (B); Heart shape development of somatic embryo (C); Heart shape somatic embryo (D); Torpedo stages of somatic embryo (red arrow) (E); and Cotyledonary stages of somatic embryos (blue arrow) in liquid cell suspension culture (F), (Bar=500µm)	5.34
5.23.	Maintenance of cell suspension culture as stock material for continuous production of somatic embryos. Bars = 1cm	5.35
5.24.	Maturation of somatic embryos derived from solid culture medium on hormone free MS medium (Bar=5mm)	5.38
5.25.	Maturation of somatic embryos derived from cell suspension culture on hormone free MS medium (Bar=5mm)	5.38
5.26.	A clump of somatic embryos (consisting of 10 to 15 of somatic embryos) which were produced from solid culture system cultured onto different concentrations of GA3 for germination (Bar=1mm)	5.39
5.27.	A clump of somatic embryos (consisting of 10 to 15 of somatic embryos) which were produced from cell suspension culture system was cultured onto different concentrations of GA3 for germination (Bar=1mm)	5.39
5.28.	Somatic embryos which were produced from solid culture system started to germinate on hormone free germination medium at the sixth to the seventh week of culture (Bar=5mm)	5.40
5.29.	Somatic embryos which were produced from cell suspension culture technique started to germinate on hormone free germination medium at the fourth to the fifth week of culture (Bar=5mm)	5.40
5.30.	Somatic embryos produced from solid culture system developed into plantlets on hormone free germination medium at the tenth week of culture (Bar=1cm)	5.41

- 5.31. Somatic embryos produced from cell suspension culture system developed into plantlets on hormone free germination medium at the eighth week of culture (Bar=1cm) 5.41
- 5.32. Normal plantlets produced from somatic embryos derived from solid culture (A), and cell suspension culture systems (B) germinated on hormone free MS medium after 12th weeks of culture (Bar=1cm) 5.42
- 5.33. Abnormal plantlets produced from somatic embryos derived from solid culture with thickness of leaves and no root was produced (A), and cell suspension culture systems with abnormal shape, stunted growth and somatic embryo growing like a vase (red arrow) (B) germinated on germination medium containing 1.5 mgL⁻¹ GA₃ after 12th weeks of culture (Bar=8mm) 5.43